

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 31/557		A1	(11) International Publication Number: WO 00/40248
			(43) International Publication Date: 13 July 2000 (13.07.00)
(21) International Application Number: PCT/SE00/00006			(81) Designated States: AE, <u>AL</u> , AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
(22) International Filing Date: 4 January 2000 (04.01.00)			
(30) Priority Data: 9900025-9 8 January 1999 (08.01.99) SE			
(71) Applicant (for all designated States except US): SYNPHORA AB [SE/SE]; Dag Hammarskjölds väg 10A, S-751 83 Uppsala (SE).			
(72) Inventors; and (75) Inventors/Applicants (for US only): STJERNSCHANTZ, Johan [FI/SE]; Villavägen 1 B, S-752 38 Uppsala (SE). RESUL, Bahram [SE/SE]; Vitkålsgratan 112, S-754 49 Uppsala (SE).			
(74) Agents: HOLMBERG, Martin et al.; Bergenstråhle & Lindvall AB, P.O. Box 17704, S-118 93 Stockholm (SE).			
(54) Title: METHOD AND COMPOSITION FOR TREATMENT OF FEMALE SEXUAL DYSFUNCTION			
(57) Abstract <p>The present invention makes available a composition for the treatment of female sexual dysfunction, said composition comprising a selective EP₂ or EP₄ prostanoid receptor agonist and in particular one of 19-hydroxy-PGE₂, 19R-hydroxy-PGE₂, 18-hydroxy-PGE₂, 20-hydroxy-PGE₂, or salts or esters thereof. The invention further discloses a method of treatment of female sexual dysfunction, wherein the above composition is applied topically in the form of a cream, gel, ointment or solids.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Larvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

METHOD AND COMPOSITION FOR TREATMENT OF FEMALE SEXUAL DYSFUNCTION

Field of invention

The present invention relates to the treatment of female sexual dysfunction, and more particularly to a novel prostaglandin based composition therefore and the use thereof for treating female sexual dysfunction.

Background of the invention

Female sexual dysfunction is not a well defined clinical entity, but can be characterized as the equivalence of male erectile dysfunction i.e. the inability of the female to reach sexual satisfaction. In contrast to the mechanism and etiology of male erectile dysfunction which have been extensively studied, only little is known about female sexual dysfunction; however, a reasonable hypothesis is inadequate vasodilatory capacity of the external genitals. Thus local drugs that restore the vasodilatory capacity of the genitals to the appropriate stimuli can be anticipated to improve the female sexual function. The prevalence of female sexual dysfunction is largely unknown, and dependent on age, but generally it can be estimated that such dysfunction is common e.g. amongst post menopausal women. Currently there is no adequate treatment of female sexual dysfunction other than hormonal treatment in post menopausal women.

While it is understood that many drugs with vasodilatory capacity administered e.g. orally may have effect on the genitals, and that certain systemic drugs e.g. sildenafil used for male erectile dysfunction, may have beneficial effect also for the treatment of female sexual dysfunction, it is a clear advantage if the drugs can be administered locally in the genital tract to avoid systemic side-effects.

Summary of the invention

We have now unexpectedly found that certain prostaglandin analogues of the E-type, more specifically agonists of the EP₂ prostanoid receptor, exert good relaxant effect in the penile cavernous tissue, and in blood vessels while exhibiting markedly reduced pain inducing effect typical for many prostaglandins. Since the female external genitals embryologically correspond to the male genitals such as the penis, and expansive cavernous tissue is present in the genitals of both sexes it is very likely that the effect of certain prostaglandin analogues is fundamentally similar in the male and female. Such drugs may therefore also be used to treat female sexual dysfunction.

In particular we have found that PGE₂ analogues substituted in carbon 18, 19 or 20 with hydroxyl (OH) exert beneficial effect, and especially one analogue, namely 19R-OH-PGE₂, exhibited a surprisingly advantageous effect with respect to relaxation of cavernous tissue and blood vessels without inducing pain as studied in an animal model.

Thus, in a first aspect, the present invention provides a composition for the treatment of female sexual dysfunction, comprising a prostaglandin which is a selective EP₂ or EP₄ receptor agonist, or an active derivative, including the free acid, or a salt or an ester thereof, optionally together with a physiologically acceptable carrier or vehicle.

In a second aspect, the invention provides a method of treating female sexual dysfunction, which method comprises administering a therapeutically active and physiologically acceptable amount of the composition to an individual in need thereof.

In a third aspect, the invention provides the use of a selective EP₂ or EP₄ receptor agonist, or an active derivative including the free acid, a salt or an ester thereof, for the preparation of a medicament for the treatment of female sexual dysfunction.

Brief description of the Figures

Fig. 1 is a diagram showing the concentration-response relation for PGE₁ and 19R-hydroxy-PGE₂ in isolated preparation of human corpus cavernosum precontracted with 10⁻⁶ M norepinephrine. Fig. 2 is a diagram showing the vasodilatory effect of PGE₁ and


19R-hydroxy-PGE₂ on rabbit penile vasculature precontracted with 10⁻⁶ M norepinephrine.

Detailed description of the invention


Engorgement of the genital tissue is primarily based on marked relaxation of both arteries and cavernous tissue as demonstrated e.g. in the penis. PGE₁ and the EP₂ receptor agonists cause vasodilation and relax the expansive tissue to about the same extent and thus promote an erectile response. However, PGE₁ being a natural prostaglandin with low selectivity for a specific prostanoid receptor has a marked irritant effect and causes pain as evident in clinical trials performed with alprostadil e.g. Padha-Natham, 1997) and as will also be shown in the experimental part further below. According to the invention, however, prostaglandin analogues with selectivity primarily for the EP₂ receptor which induce the same beneficial effect but with significantly less irritant effect are much more preferable as it can be predicted that such analogues will cause no or only minimal pain in clinical practice. To facilitate the understanding of the invention, a general description of prostaglandins is first given.

The prostaglandins are fatty acids usually derived from the precursors eicosatrienoic, eicosatetraenoic or eicosapentanoic acid through metabolic steps involving oxygenation. The prostaglandins typically carry a cyclopentane ring to which two carbon chains link, the upper chain usually being called the alpha chain and the lower the omega chain. The alpha chain is a 7 carbon chain with a terminal carboxylic acid moiety whereas the omega chain is an 8 carbon methyl-terminated aliphatic chain. Depending on the number of double bonds in these chains subscripts of 1-3 are used. In prostaglandins with subscript 1, e.g. PGE₁, the double bond is situated between carbons 13 and 14 in the omega chain, and it exhibits trans configuration in naturally occurring prostaglandins. In prostaglandins with subscript 2, e.g. PGE₂, an


additional double bond in the *cis* configuration exists between carbons 5 and 6 in the alpha chain, and finally in prostaglandins with subscript 3 a third double bond is situated between carbons 17 and 18 in the omega chain. This double bond also exhibits *cis* configuration in naturally occurring prostaglandins. All naturally occurring prostaglandins carry a hydroxyl group in carbon 15, which is essential for biological activity. The substituents and the configuration of the cyclopentane ring determine whether the prostaglandin is of the A, B, C, D, E, F, G, H, I or J type as depicted in Scheme 1 below. The prostaglandins that have been used to exemplify the present invention are of the E-type, and the chemical structures of these prostanoids are depicted in Scheme 2. Except for 18RS-OH-PGE₂-methyl ester the prostaglandin analogues were also used as acids.




Scheme 1




PGE1-Isopropyl ester



18RS-Hydroxy-PGE2-methyl ester



19R-Hydroxy-PGE2-methyl ester



20-Hydroxy-PGE2-methyl ester

Scheme 2

The prostaglandins exert their pharmacologic effect through specific G protein coupled membrane receptors. There are at least 8 different receptors for the endogenous prostaglandins, namely the FP ($\text{PGF}_{2\alpha}$), EP_1 , EP_2 , EP_3 , EP_4 (PGE_2), DP (PGD_2), IP (PGI_2), and TP (TxA_2) receptors, the most common naturally occurring ligand for respective receptor being shown within parentheses. At least for the EP_3 receptor splice variants have been shown to exist. The EP_2 and EP_4 receptors are closely related, and probably mediate similar effects. We have therefore studied the effect only on the EP_2 receptor, and regard it to represent the EP_4 receptors as well in the case the compounds studied would have an effect on that receptor, too. The

molecular biology and pharmacology of the prostanoid receptors have recently been reviewed by Coleman et al., 1994.

From a therapeutic point of view a problem with the endogenous prostaglandins is that they exert effects on many different prostanoid receptors. Each endogenous prostaglandin has a preference for one specific receptor type, but is not very selective and usually distinguishes poorly between the receptor subtypes, i.e. the EP receptors. Thus PGE₁ and PGE₂ are good ligands for all subtypes of the EP receptor. Consequently selective effects on one of the subtypes of the EP receptor is impossible to achieve with the endogenous prostaglandins. However, certain synthetic prostaglandin analogues, e.g. butaprost, 11-deoxy-PGE₁ and AH13205 as well as a naturally occurring metabolite of PGE₂, namely 19R-OH-PGE₂, are selective EP₂ prostanoid receptor agonists.

18-OH-PGE₂, 19R-OH-PGE₂ and 20-OH-PGE₂ are effective EP₂ receptor agonists with selectivity for the EP₂ receptor over the EP₃ receptor. The endogenous PGE₁ is unselective and does not distinguish between the EP receptor subtypes sufficiently, and furthermore significantly spills over on the DP/IP receptors which e.g. the 18-, 19- and 20-OH substituted PGE₂ analogues do not. However, PGE₁ has been included as a reference substance as it is the only prostaglandin currently in clinical use for the treatment of erectile dysfunction.

Accordingly, high selectivity or specificity to the EP₂ receptor compared to other prostaglandin receptors, particularly the EP₃ receptor, characterizes the compounds to be used in the method or compositions according to the present invention. It need not be said that the more specific the compound is for the EP₂ receptor the better results are obtained, but a certain advantage is, of course, achieved also in cases of some interaction with other receptors. High selectivity in this connection means that the effect on the EP₂ receptor is at least more than 5 times, especially more than 10 times, and in particular more than 100 or 1000 times the effect on other prostaglandin receptors.


As stated above, according to the present invention, in particular the selective EP₂ receptor agonist 19R-OH-PGE₂ and its carboxylic acid esters appear to be unique and ideally suited for the treatment of female sexual dysfunction. In this context it should be mentioned that 19R-OH-PGE₂ is a metabolite of PGE₂ in the genital tract and can be found in large quantities normally in human semen (Taylor and Kelly, 1974).

However, the physiological role of this unique metabolite is unknown. Thus, 19R-OH-PGE₂ and its carboxylic acid esters constitute a very attractive medication for female sexual dysfunction since this analogue has a marked vasodilatory capacity, causes no pain, and furthermore occurs normally in the body.

With respect to 19R-OH-PGE₂ which appears ideally suited for treatment of female sexual dysfunction, it should be observed that various modifications or substitutions of the molecule are possible as long as the new derivatives exhibit selective agonism on the EP₂ receptor.

In Formula I, the wavy bonds represent the α (S) or β (R) configuration, and the dashed bonds represent a single, a triple or a double bond in the cis or trans configuration. R in the formula is hydrogen, a salt moiety, e.g. an alkali or ammonium salt, a straight or branched saturated or unsaturated alkyl group, preferably with 1-10 carbons, an alicyclic ring, preferably with 3-8 carbons, arylalkyl, preferably aryl-C₂₋₅ alkyl, or an aryl ring. X is a straight chain, saturated or unsaturated, preferably consisting of 2-5 carbons optionally interrupted by a heteroatom (O, S, N), and the chain may contain an alicyclic ring, preferably C₃-C₇ cycloalkyl, or an aryl or heteroaryl ring. R₁ and R₂, which are identical or different, are hydrogen, hydroxy, halogen, oxygen (=O or alkoxy) or an alkyl group with 1-3 carbons or an alkoxy group with 1-3 carbons or an ester OCOR₃, where R₃ is a straight or branched, saturated or unsaturated alkyl group preferably containing 1-10, especially 1-6 carbons, or a cycloalkyl, preferably containing 3-7 carbons, or an aryl or arylalkyl group, preferably aryl-C₂₋₅ alkyl (e.g. benzyl). Optionally C₁₀ can be substituted e.g. to contain a mono- or dialkyl group.

Z is an alkyl chain of 1-8, e.g. 3-8 and 2-4 carbons, saturated or unsaturated, optionally interrupted by one or more heteroatoms (O, S, N), and containing one or more, preferably one substituent Y, attached to a carbon atom within the chain or a carbon atom in terminal position. The chain Z may be branched, containing alkyl (preferably methyl) substituents and/or contain an alicyclic ring, e.g. a cycloalkyl, or an aryl (esp. a benzene) or heteroaryl ring, in inter- or terminal position. Y, is hydroxy, sulfhydryl, amino, methylamino, dimethylamino, C₁₋₃ alkoxy or halogen (Cl, Br, F) or oxygen (keto). In a preferred embodiment of the invention Y is attached to carbon 18, 19 or 20. In the most preferred embodiment Y is OH and attached to carbon 19. At present it is believed that the inventive concept is based on the electronegative effect introduced by Y.



The prostaglandins may be epimeric mixtures as well as in the form of the individual epimers.

Description of suitable embodiments

The EP₂ prostanoid receptor agonists according to the present invention can be used as the normal carboxylic acids, salts (e.g. cationic) or as ester prodrugs, preferentially as carboxylic acid esters. The active compound can be administered topically in a pharmaceutically acceptable delivery medium. Cyclodextrins may be employed for solubilisation and stabilisation. Different delivery media, e.g. liniment, creams, gels, ointments, and possibly solid state forms may be employed. Gels, creams, ointments and solid state formulations may or may not contain preservative such as benzalkonium

chloride, chlorhexidine, thiomersal, parabenoic acid, and other compounds with satisfactory antimicrobial effect. The dose interval is 0.001-1 mg, typically 0.01-0.1 mg per application.

Accordingly the new medicament should be administered locally in the genital tract. Such treatment should be initiated 5-60 min before intercourse.

Exemplification of the invention

The test compounds PGE₁-isopropyl ester, 18RS-hydroxy-PGE₂-methyl ester, 19R-hydroxy-PGE₂-methyl ester and 20-hydroxy-PGE₂-methyl ester were dissolved in 0.5% polysorbate-80 as a stock solution, and were diluted in 0.5% polysorbate-80 to the appropriate concentrations in the various pharmacological experiments.

Synthesis of PGE₁-isopropyl ester

PGE₁ was obtained from Chinoin, Pharmaceutical and Chemical Works Co. Ltd., Budapest, Hungary.

DBU (644 mg, 4.2 mmol) in acetonitrile (1 ml) was added dropwise to a stirred solution of PGE₁ (300 mg, 0.84 mmol) in acetonitrile (3 ml) at 0°C. The mixture was allowed to warm to room temperature whereupon isopropyl iodide (1142 mg,

6.72 mmol) in acetonitrile (2ml) was added dropwise. After being stirred for 10h (TLC monitoring), the reaction mixture was quenched with water (8ml), the mixture was extracted with ethyl acetate (2x50ml), and the extract was washed with brine (10ml), citric acid 3% (10ml), and finally sodium hydrogen carbonate 5% (2x10ml). After drying with anhydrous sodium sulphate, the solvent was removed in vacuo and the residual oil was chromatographed on silicagel using ethyl acetate as eluent. This afforded 230 mg of the product (69%) of the title compound as a colorless oil: R_f = 0.18 (ethyl acetate); ^1H NMR (CDCl_3) δ 0.89 (m, 3H), 1.2 (d, 6H), 1.21-1.4 (m, 10H), 1.42-1.62 (dm, 6H), 2.2-2.4 (dm, 4H), 2.7-2.75 (dd, 1H), 4.0-4.17 (dm, 2H), 5.0 (m, 1H), 5.5-5.7 (dm, 2H).

Synthesis of 18RS-hydroxy-PGE₂-methyl ester

1. Preparation of Dimethyl-(2-oxo-5-heptyne)-phosphonate.(2)

To a stirred suspension of dry sodium hydride (3.13 g, 124,07 mmol) in dry THF (100 ml) at room temperature was added dropwise a solution of dimethyl-(2-oxopropyl)-phosphonate (20,61 g, 124,07 mmol) in dry THF (50 ml) under N_2 . The reaction mixture was stirred for 1h, then cooled in an ice-bath and treated with a solution of n-butyllithium (7.95 g, 124, 07 mmol) in hexane, causing a dark brown solution to be formed. Stirring was continued for 1 h at 0 °C, followed by dropwise addition of 1-bromo-2-butyne.1 (15 g, 112,79 mmol) in THF (30 ml). The reaction mixture was gradually warmed to room temperature and after 1 h (TLC monitoring) the reaction mixture was quenched with ice-water, HCl 1M to pH about 4 and extracted, twice, with ethyl acetate. The organic layer was washed with brine and chromatographed on silica gel using EtOAc as eluent. R_f =0.38 (silica gel, ethylacetate), yield 18g (73%).

2. (1S,5R,6R,7R)-6-Formyl-7-((4-phenylbenzoyl)oxy)-2-oxabicyclo {3.3.0}octan-3-one.(3)

A mixture of Coreys lactone (23,06 g, 65,44 mmol), DCC (40,50 g, 196,31 mmol), DMSO (27,8 ml, 392,6 mmol) and phosphoric acid 85% (2,2 ml) in DME (130 ml) was stirred at room temperature for 2h (TLC monitoring). The precipitate was removed on silica-gel pad washed with DME (2 x 50 ml). The filtrate was concentrated in vacuo and used for the next step without isolation.

4. (1S,5R,6R,7R)-6-(3-Oxo-6-yn-1E-1-octenyl)-7-((4-phenylbenzoyl)oxy)-2-oxabicyclo{3.3.0}octan-3-one.(4)

To a stirred suspension of sodium hydride (1,98 g, 78,53 mmol) in DME (140 ml) under N₂ was added dropwise the above phosphonate 2 (18,56 g, 85,07 mmol) and stirred mechanically at room temperature for 1,5 h. The mixture was then cooled to -5 °C, and a solution of the above crude Coreys aldehyde 3 was added dropwise. After 30 min at 0 °C and 2 h at room temperature (TLC monitoring), the reaction mixture was neutralised with citric acid 5% and extracted with ethyl acetate (2x100ml). The organic layer was dried and evaporated. The residue (oil) was chromatographed on silica-gel using ethyl acetate and 10% methanol in ethyl acetate successively, giving a light yellowish oil. R_f = 0.58(silica-gel, ethyl acetate), yield 52%.

5. (1S,5R,6R,7R)-6-(3,6-Dioxo-1E-1-octenyl)-7-((4-phenylbenzoyl)oxy)-2-oxabicyclo{3.3.0}octan-3-one.(5)

To the acetylenic solution 4 in acetonitrile:water 2:1 (100 ml) was added mercuric oxide (13,8 g, 63,73 mmol) and sulfuric acid 1M (25,49 ml, 25,49 mmol). The reaction mixture was stirred magnetically. After about 1 h at room temperature (TLC monitoring) the reaction mixture was worked-up by addition of ethylacetate and HCl 1M. The organic layer was dried and evaporated. The crude oil was used for the next step without purification. R_f = 0.44 (silicagel, EtOAc), yield = 41%.

6. (1S,5R,6R,7R)-6-(3RS,6RS-Dihydroxy-1E-1-octenyl)-7-((4-phenylbenzoyl)oxy)-2-oxabicyclo{3.3.0}octan-3-one.(6)

To a stirred solution of the above diketone 5 (12,0 g, 26.1 mmol) and cerium chloride hepta hydrate (5.83 g, 15,64 mmol) at -20 °C in methanol: methylene chlorider 1:1 was added sodium borohydride (1.48 g, 39,09 mmol) in small portions under N₂. After 30 min (TLC monitoring). The reaction mixture was quenched with HCl 1M to pH about 4-5, and diluted with water (50 ml) and ethyl acetate (100 ml). The organic layer was separated and the water layer was washed twice with EtOAc, dried and evaporated. The residue was purified on silica gel using EtOAc as eluent. The title

compound **6** was obtained as colorless oil: yield 8.8 g (71%). $R_f = 0,15, 0,13$ corresponding to the two isomers 15α and 15β (silicagel, EtOAc).

^1H NMR (CDCl_3) δ 0,9 (m, 3H), 1,4-1,8 (dm, 6H), 2,3 (d, 1H), 2,5-2,9 (dm, 5H), 3,5 (m, 1H), 4,2 (m, 1H), 5,1 (m, 1H), 5,3 (m, 1H), 5,7 (m, 2H), 7,4 (m, 1H), 7,5 (m, 2H), 7,6-7,7 (dd, 4H), 8,1 (d, 2H); ^{13}C NMR (CDCl_3) δ 176,40 ($\text{C}_6\text{H}_4\text{C}=\text{O}$), 16591 (lactone $\text{C}=\text{O}$), 146,07, 139,83, 136,21, 130,15, 128,91, 128,31, 127,15, 83,27, 79,13, 73,11, 71,79, 71,48, 54,08, 42,84, 42,75, 37,55, 34,85, 34,01, 33,43, 32,95, 32,09, 30,17, 9,96.

7. (1S,5R,6R,7R)-6-(3RS,6RS-Di t butyldimethylsilyloxy-1E-1-octenyl)-7-((4-phenylbenzoyl)oxy)-2-oxabicyclo{3.3.0}octan-3-one.(7)

To a stirred solution of the above dihydroxy compound **6** (8,6 g, 18,51 mmol) in dichloromethane was added triethyl amine (12,83 ml, 92,56 mmol), t-butyldimethylsilyl chloride (13,95 g, 92,56 mmol) and 4-dimethylamino pyridine (1,13 g, 9,26 mmol). The mixture was stirred magnetically for 15 h at room temperature. The reaction mixture was diluted with ether, filtered and the precipitate was washed with ether. The organic layer was washed with brine, dried and concentrated in vacuo. The residue was chromatographed on silica gel using 5% ether in dichloromethane. $R_f = 0.58$ (silica gel, 5% ether in CH_2Cl_2). Yield 12,2 g (92%)

8. (1S,5R,6R,7R)-6-(3RS,6RS-Di t butyldimethylsilyloxy-1E-1-octenyl)-7-((4-hydroxy)-2-oxabicyclo{3.3.0}octan-3-one.(8)

To a stirred solution of the above disilyl ether **7** (12 g, 17,31 mmol) in methanol was added potassium carbonate (1,2 g, 8,66 mmol). The reaction mixture was stirred at room temperature for 4 h (TLC monitoring). The mixture was neutralised with citric acid 5%, extracted twice with ethyl acetate (100 ml), dried and concentrated in vacuo. The oil was chromatographed on silica gel using gradient elution with 5% ether in CH_2Cl_2 and EtOAc:acetone 1:1 successively. $R_f = 0.43$ (silica gel, ethyl acetate), yield = 8,86 g (74%)

9. (1S,5R,6R,7R)-6-(3RS,6RS-Di t butyldimethylsilyloxy-1E-1-octenyl)-7-((4-t-butyldimethylsilyloxy)-2-oxabicyclo{3.3.0}octan-3-one.(9)

To a stirred solution of the above disilyloxy ether 8 (6,6 g, 12,86 mmol) in CH_2Cl_2 at room temperature was added triethyl amine (7,14 ml, 51,47 mmol) t-butyldimethylsilyl chloride (7,76 g, 51,47 mmol) and 4-dimethyl aminopyridine (0,47 g, 3,9 mmol). The reaction mixture was worked-up as in 7. The crude product was chromatographed on silica gel using 5% ether in dichloromethane to give a pure trisilyloxy ether product 9 as an oil. $R_f = 0.62$ (silica gel, 5% ether in CH_2Cl_2), yield 7,8 g, (96%)

10. (1S,5R,6R,7R)-6-(3RS,6RS-Di t butyldimethylsilyloxy-1E-1-octenyl)-7-((4-t butyldimethylsilyloxy)-2-oxabicyclo{3.3.0}octan-3-ol.(10)

A solution of diisobutyl aluminium hydride (DIBAL-H) (2,9 g, 20,87 mmol) in dry THF was added dropwise to a stirred solution of the above trisilyl ether lactone 9 (7,7 g, 12,24mmol) in THF (80 ml) at -72/-80 °C. After 1h (TLC monitoring), the reaction mixture was quenched with ice (15 g) and ethyl acetate (150 ml), filtered and the filtrate was concentrated in vacuo. The residue was used directly without separation in the next step. $R_f = 0.27$ (Silica gel, 5% ether in dichloromethane).

11. 11,15RS,18RS-Tri-t-butyldimethylsilyloxy-PGF_{2α} (11)

To a stirred suspension of 4-carboxybutyl-triphenylphosphonium bromide (21,70 g, 48,95 mmol) in dry THF under N_2 at -5 °C was added potassium tert-butoxide (10,99 g, 97,91 mmol) and the mixture was stirred at room temperature for 30 min. To the resultant red-orange solution of ylide at -15/-10 °C was added the lactol 10 (7,7 g, 12,24 mmol) in THF, and the mixture was stirred for 2-3 h (TLC monitoring) at room temperature. The reaction mixture was diluted with water and quenched with citric acid 15 % to pH 6-7 and extracted with EtOAc, dried and concentrated in vacuo. The resultant slurry was used directly without isolation for the next step.

12. 11,15,18RS-Tri-t-butyldimethylsilyloxy-PGF_{2α} methyl ester (12)

Methyl iodide (8,6 g, 61,20 mmol), was added to a stirred solution of the crude product 10 (8,73 g, 12,24 mmol) and N,N-Diisopropyl ethyl amine (9,473 g, 73,44 mmol) in acetonitrile at room temperature. After 15h (TLC monitoring) the mixture was diluted with water (100 ml) and ethyl acetate (150 ml), washed with sodium hydrogen carbonate 5% (60 ml) and brine (70 ml). The organic layer was dried and

evaporated in vacuo. The residue was chromatographed on silica gel using EtOAc as eluent. $R_f = 0.18$ (silica gel, EtOAc:hexane 1:9).

13. 11,15,18RS-Tri-t-butyldimethylsilyloxy-PGE₂ methyl ester (13)

To a stirred solution of the compound 11 (3,3 g, 4,54 mmol) in dichloromethane was added pyridinium chlorochromate (PCC) (3,9 g, 18,15 mmol) treated with aluminium oxide (1g PCC was stirred with 5g aluminium oxide in acetone, the solvent was removed in vacuo giving a light yellow powder). The resulting suspension was stirred at room temperature for 4h (TLC monitoring). The suspension was filtered on silica gel pad, washed with dichloromethane. The solvent was removed and the resulting oil was diluted with ether and washed with water (50 ml), sodium hydrogen carbonate 5% (50 ml). The solvent was removed in vacuo. The residue was chromatographed on silica gel using 5% ether in CH₂Cl₂. $R_f = 0.32$ (silica gel, EtOAc:hexane 1:9), yield 3,1 g (86%)


14. 18RS-hydroxy PGE₂ methyl ester (14)

The protecting groups 11,15,18-tri-t-butyldimethylsilyl chloride were removed by addition of HF 4% (108 ml) to a solution of 13 (3,0 g, 3,98 mmol) in acetonitrile (300 ml). The reaction mixture was stirred at room temperature for about 8h (TLC monitoring). The reaction mixture was worked-up by addition of EtOAc (200 ml). The organic layer was separated and washed with sodium bicarbonate 5% and the pH was adjusted to about 6. The organic layer was washed with brine, dried and concentrated in vacuo. The residue was purified by column chromatography on silica gel using gradient elution with CH₂Cl₂, EtOAc and 5-10% methanol in ether successively (The stationary phase; silica gel, in the column must be washed with the eluent containing triethylamine before purification, to avoid isomerization) 18RS-hydroxy-15S-PGE₂ Me ester $R_f = 0.16$ (5% MeOH in ether); yield 310 mg.

18RS-hydroxy-15R-PGE₂ Me ester $R_f = 0.20$ (5% MeOH in ether);
yield 248 mg.

18RS-hydroxy-PGE₂ methyl ester ¹H NMR (CDCl₃) δ 0,9 (t, 3H), 1,4-1,7 (dm, 8H), 2,1 (m, 2H), 2,2 (m, 2H), 2,3-2,4 (m, 5H), 2,7 (m, 1H), 3,6 (m, 1H, 18-CHOH), 3,7 (s, 3H), 4,05, (m, 1H, 15-CHOH), 4,2 (m, 1H, 11-CHOH), 5,3-5,5 (dm, 2H, db), 5,6-5,8 (dm, 2H, db);

^{13}C (NMR) δ 214, 174, 24, 136, 7, 131, 3, 130, 8, 126, 5, 72, 36, 71, 78, 71, 70, 55, 54, 54, 54, 53, 70, 51, 60, 46, 06, 34, 06, 33, 42, 33, 00, 32, 10, 30, 34, 30, 29, 30, 07.



Scheme 3

Synthesis of 19-R-hydroxy-prostaglandin E_2 -methyl ester

19R-hydroxy-prostaglandin E_2 was obtained from Cayman Chemicals, Ann Arbor, Michigan, USA. Methyl iodide (9.2 mg, 0.065 mmol) in acetonitrile (1.0 ml) was added drop-wise to a stirred solution of 19R-hydroxy-prostaglandin E_2 (4 mg, 0.011 mmol), and N,N-diisopropyl ethyl amine (7 mg, 0.054 mmol) in acetonitrile. More methyl iodide (4.5 mg, 0.032 mmol) in acetonitrile was added after 6 h and the stirring was continued for 12 h (TLC monitoring). The reaction mixture was quenched with water (5.0 ml) and extracted with ethyl acetate (2x10 ml), and the organic phase was washed with sodium hydrogen carbonate 5% (5 ml) and hydrochloric acid 0.5M (5 ml). After drying with anhydrous sodium sulphate, the solvent was removed in vacuo and the residue was chromatographed

on silica-gel using ethyl acetate: acetone 1:1 and acetone as eluent. This afforded 3.2 mg (72.7%) of the product as a colorless oil: $R_f=0.17$ (ethyl acetate:acetone:acetic acid 1:1:0.01) $^1\text{H NMR}$ (CDCl_3) δ 1.25 (d, 3H), 1.5-1.7 (m, 8H), 2.1-2.6 (m, 9H), 3.7 (s, 3H), 3.8 (m, 1H), 4.1 (m, 1H), 4.2 (m, 1H), 5.3-5.5 (dm, 2H), 5.6-5.8 (dm, 2H).

Synthesis of 20-OH-prostaglandin E_2 -methyl ester

The commercially available 20-hydroxy PGE_2 (Cayman Chemicals, Ann Arbor, Michigan) (2.0 mg, 0.0054 mmol) was esterified in acetonitrile (2.0 ml) with methyl iodide (4.6 mg, 0.0327 mmol) in the presence of N,N -diisopropyl ethylamine (3.5 mg, 0.027 mmol). The reaction mixture was stirred at room temperature for 10 h (TLC monitoring, silica gel, ethyl acetate). The reaction mixture was quenched with water (3.0 ml) and extracted with ethyl acetate (2x10 ml). The organic layer was dried and concentrated in vacuo and the residual oil was chromatographed on silica gel using ethyl acetate:acetone 1:0.5 as eluent; $R_f = 0.38$ (silica gel, ethyl acetate:acetone 1:1).

Pharmacological experiments

The effect of the test compounds on cavernous tissue was tested using penile tissue. As mentioned previously both in the male and female the genitals contain cavernous tissue, and the penile tissue can therefore be used as model for genital cavernous tissue. The general blood vessel relaxing effect of the test compounds was studied by investigating the immediate blood pressure decrease in rats upon intravenous infusion of the test compounds. The acute drop in blood pressure reflects a peripheral vasodilation in the animals brought about by the test compounds. The general irritative effect of the test compounds was studied in the cat eye which is very sensitive to noxious stimuli.

Human penile cavernous tissue was obtained fresh from surgery, and representative tissue samples were mounted in smooth muscle tissue baths containing a modified Krebs' solution consisting of NaCl 119 mM, KCl 4.6 mM, MgCl_2 1.2 mM, NaH_2PO_4 1.2 mM, NaHCO_3 15 mM, CaCl_2 1.5 mM and glucose 11 mM. The solution also contained indomethacin at a final concentration of about $3\mu\text{M}$. The solution was continuously bubbled by 95% O_2 and 5% CO_2 , and the temperature was kept at 37°C .

The tissue preparations were stretched by a force corresponding to 500 mg, and were given a contractile tone by the addition of norepinephrine at a concentration of 10^{-6} M. Concentration-response curves were then constructed by adding cumulatively increasing concentrations of the prostaglandin analogues to the bath in a routine way. The relaxant effect was normalized by comparing to that of carbachol in the same preparation. PGE₁ and 19R-OH-PGE₂, respectively, were used as acids.

To study the effect of the PGE₁ and 19R-OH-PGE₂ on the penile vasculature, penile blood vessels of the rabbit were isolated and mounted as ring segments in a small vessel myograph (J.P. Trading, Denmark) containing a solution consisting of NaCl 119 mM, KCl 4.7 mM, CaCl₂ 1.5 mM, MgSO₄ 1.17 mM, KH₂PO₄ 1.18 mM, NaHCO₃ 25 mM, EDTA 0.027 mM and glucose 11 mM. The solution also contained indomethacin at a final concentration of about 3 μ M, and was continuously bubbled by 95% O₂ and 5% CO₂. The temperature was kept constant at 37 °C. The vessels were stretched and then precontracted using 10^{-6} M norepinephrine, and the vasorelaxant effect of the prostaglandin analogues was normalized by comparing it to that of papaverine in the same preparation. Cumulative concentration-response curves were constructed for the analogues.

The vasodilatory effect of the prostaglandin analogues was also studied in the rat by registering the blood pressure reducing effect. This is a relevant in vitro model for showing general vasodilatory effect of importance for inducing erection since vasodilation in penis is needed for achieving erection. Rats were anaesthetized with a mixture of ketamine and xylazin and the prostaglandin analogues were infused intravenously. The blood pressure was continuously registered in a femoral artery. Each analogue was infused in 3 escalating doses in the same animal. None of the compounds was found to have any significant effect of the heart rate and the immediate reduction in arterial blood pressure therefore reflects an acute vasodilatory response to the prostaglandin analogues tested. In all experiments the blood pressure reduction was transient and the blood pressure increased immediately after termination of the infusion. The analogues were used both as esters and acids for the experiments.

The irritant effect of the prostaglandin analogues was tested using a behavioural model in cats. In this model the irritant effect on the eye is studied by registering the degree of eyelid closure and the behaviour of the animals. Compounds that cause discomfort and pain in the eye cause the animals to close their eyes. The analogues were applied as carboxylic acid esters, to enhance bioavailability, topically on the eye as a single dose at different dose levels. The animals were then followed for several hours at regular intervals. Each compound and dose was tested on a group of 3-6 cats. At least 3 days elapsed between two consecutive tests on the same animals. An arbitrary scale from 0 (absence of irritation) to 3 (marked irritation), with half steps was used.

The relaxant effect on the human penile cavernous tissue of PGE_1 and 19R-OH-PGE_2 are presented in Table I. It can be seen that PGE_1 and 19R-OH-PGE_2 were about equipotent and comparable to carbachol, however PGE_1 was slightly more potent than 19R-OH-PGE_2 . Since 19R-OH-PGE_2 is a selective EP_2 receptor agonist this finding indicates that the EP_2 receptor accounts for most of the relaxant effect of PGE_1 . Furthermore, in Fig.1 it is demonstrated that the concentration-response curves of PGE_1 and 19R-OH-PGE_2 are parallel and differ only with a factor of about 2-3, i.e. the latter analogue is about half to one third as potent as the former.

Table I. Relaxation of human corpus cavernosum tissue induced by prostaglandins compared to carbachol (10^{-6} M). (Mean \pm SEM)

Compound	n	Reduction in tension (%)	EC-50 (Moles/l)
Carbachol	8	100.0 \pm 0.0	—
PGE_1	8	90.9 \pm 3.7	2.5×10^{-7}
19R-OH-PGE_2	8	96.9 \pm 3.1	6.9×10^{-7}

The vasodilatory effect of PGE_1 and the EP_2 receptor agonist 19R-OH-PGE_2 on rabbit penile blood vessels is demonstrated in Fig 2. It can be seen that the two

prostaglandins had the same efficacy in inducing vasodilatation and were also about equipotent. This furthermore demonstrates that the EP₂ receptor mediates most of the vasodilatory effect of PGE₁. The vasodilatory effect of all the prostaglandin analogues as studied in the rat by investigating the immediate reduction in blood pressure upon intravenous infusion is presented in Table II. As can be seen in the table PGE₁ and the three hydroxy-substituted PGE₂ analogues effectively reduced the blood pressure in the anaesthetized rat, demonstrating an acute vasodilatory response.

Table II. Reduction of blood pressure in anaesthetized rats by prostaglandin analogues with agonistic effect on the EP₂ receptor. (n= 3 for each dose; Mean±SEM)

Prostaglandin analogue	2.5 - 3.5 µg/kg		7.5 - 10 µg/kg		25 - 35 µg/kg	
	Blood pressure o (mmHg)	Reduction (%)	Blood pressure o (mmHg)	Reduction (%)	Blood pressure o (mmHg)	Reduction (%)
PGE ₁	114±20	13±4	105±15	35±5	103±9	52±3
18RS-OH-PGE ₂	122±28	7±3	123±33	15±3	115±32	32±2
19R-OH-PGE ₂	115±14	13±10	103±12	31±3	91±9	43±6
20-OH-PGE ₂	91±11	26±5	85±10	38±3	91±6	44±0

Blood pressure o = blood pressure before infusion of prostaglandin analogue.

The irritant effects of the three prostaglandin analogues and PGE₁ as studied in the feline eye are presented in Table III. All three hydroxy-substituted PGE₂-analogues had significantly less irritant effect than PGE₁-isopropyl ester and PGE₁ acid. Most surprisingly we found that both 18RS-OH-PGE₂-methyl ester and 19R-OH-PGE₂-methyl ester had no or markedly reduced irritant effect, even at doses 1000 times higher than PGE₁-isopropyl ester. Previously it has been shown that 19R-OH-PGE₂ does not cause lid closure in rabbits but induces an irritant effect by inducing swelling of the ocular structures (Hall and Jaitly, 1977). We found no evidence of such irritation in the cat eye with 19R-OH-PGE₂ or the other hydroxy-substituted PGE₂ analogues.

To confirm that 19R-OH-PGE₂-methyl ester, which is more hydrophilic than PGE₁-isopropyl ester, indeed penetrates into the cornea and the intraocular parts of the eye we measured the intraocular pressure in 3 cats under local anaesthesia before and 1

hour after topical administration of the prostaglandin to the eye. The intraocular pressure was measured by pneumatonometry. One eye was treated with the test compound and the other eye received the vehicle only. In the eyes treated topically with 19R-OH-PGE₂-methyl ester the intraocular pressure decreased from 16.3 \pm 0.9 mmHg to 12.7 \pm 1.2 mmHg, whereas it was 16.3 \pm 0.9 mmHg and 15.7 \pm 0.7 mmHg in the control eyes at the same time points. It is well known that prostaglandins reduce the intraocular pressure in cats (Bito et al., 1989), and it can thus be taken as an evidence that the drug has penetrated into the eye. Furthermore, from Table III it can also be seen that the PGE₁ acid, which is a much less lipophilic compound than the 18-, 19- and 20-hydroxy-substituted PGE₂-methyl ester analogues, caused significantly more irritation, and at much lower dose, than the hydroxy-substituted analogues. Thus the absence of pain and irritation after administration of 18RS-OH-PGE₂, 19R-OH-PGE₂ and 20-OH-PGE₂ cannot be explained on the basis of increased hydrophilicity and poor bioavailability. Accordingly, selective EP₂ prostanoid receptor agonists seem to be very advantageous in that they exhibit no, or markedly reduced irritant effect.

Table III. Maximum irritative effect of prostaglandin analogues as studied in cat eyes.

The log P values were estimated based on thin layer chromatography with PGF_{2 α} isopropyl ester as reference (log P 4.5). ie= isopropyl ester, me= methyl ester.

Maximum irritation =3. Mean \pm SEM (n= 3-6).

Analogue	log p-value	Dose: 0.01 μ g	Dose: 0.1 μ g	Dose: 0.3 μ g	Dose: 1 μ g	Dose: 3 μ g	Dose: 10 μ g	Dose: 30 μ g	Dose: 100 μ g
PGE ₁ -ie	4.7	0.4 \pm 0.2	1.7 \pm 0.3	2.8 \pm 0.1	-	-	-	-	-
PGE ₁ acid	0.7	-	0.6 \pm 0.1	-	1.6 \pm 0.2	-	3.0 \pm 0.0	-	-
18RS-OH-PGE ₂ -me	2.7	-	-	-	0.0 \pm 0.0	0.0 \pm 0.0	0.0 \pm 0.0	0.3 \pm 0.2	-
19R-OH-PGE ₂ -me	2.5	-	0.0 \pm 0.0	-	0.1 \pm 0.1	-	0.1 \pm 0.1	-	0.7 \pm 0.4
20-OH-PGE ₂ -me	2.2	-	-	-	-	0.5 \pm 0.0	1.2 \pm 0.2	1.2 \pm 0.4	-

References

Bito, L.Z., Camras, C.B., Gum, G.G. et al. (1989). The ocular hypotensive effect and side-effects of prostaglandins on the eyes of experimental animals. In: The ocular effects of prostaglandins and other eicosanoids (Eds.: L.Z. Bito and J. Stjernschantz). Alan R. Liss Inc. New York. Pp: 349-368.

Coleman, R.A., Smith, W.L. and Narumiya, S. (1994). VIII. International Union of Pharmacology classification of prostanoid receptors: Properties, distribution and structure of the receptors and their subtypes. Pharmacological Reviews. 46; 205-229.

Hall, D.W.R. and Jaitly, K.D. (1977). Inflammatory responses of the rabbit eye to prostaglandins. Agents Actions. Suppl. 2: 123-133.

Pada-Nathan, H., Hellstrom, W.G.J., Kaiser, F.E. et al. (1997). Treatment of men with erectile dysfunction with transurethral alprostadil. New Eng. J. Med. 336; 1-7.

Taylor, P.L. and Kelly R.W. 1974. 19-hydroxylated E prostaglandins of human semen. Nature. 250; 665-667.

Claims

1. The use of a prostaglandin for the manufacture of a pharmaceutical composition for the treatment of female sexual dysfunction, **characterized** in that said prostaglandin is a selective EP₂ receptor agonist or an active prostaglandin derivative thereof, such as a salt or an ester thereof.
2. The use of a prostaglandin for the manufacture of a pharmaceutical composition for the treatment of female sexual dysfunction, **characterized** in that said prostaglandin is a selective EP₄ receptor agonist or an active prostaglandin derivative thereof, such as a salt or an ester thereof.
3. The use of 19-hydroxy-PGE₂, or a salt or ester thereof, for the manufacture of a pharmaceutical composition for the treatment of female sexual dysfunction.
4. The use of 19R-hydroxy-PGE₂, or a salt or ester thereof, for the manufacture of a pharmaceutical composition for the treatment of female sexual dysfunction.
5. The use of 18-hydroxy-PGE₂, or a salt or ester thereof, for the manufacture of a pharmaceutical composition for the treatment of female sexual dysfunction.
6. The use of 20-hydroxy-PGE₂, or a salt or ester thereof, for the manufacture of a pharmaceutical composition for the treatment of female sexual dysfunction.
7. The use of 19R-hydroxy-PGE₂-methyl ester for the manufacture of a pharmaceutical composition for the treatment of female sexual dysfunction.
8. The use of 19R-hydroxy-PGE₂-isopropyl ester for the manufacture of a pharmaceutical composition for the treatment of female sexual dysfunction.
9. A method of treating female sexual dysfunction, comprising administering to a human a therapeutically effective and physiologically acceptable amount of a prostaglandin, **characterized** in that said prostaglandin is a selective EP₂ receptor agonist.
10. The method according to claim 9, **characterized** in that said prostaglandin is chosen among 19-hydroxy-PGE₂, 19R-hydroxy-PGE₂, 18-hydroxy-PGE₂, 20-hydroxy-PGE₂, or salts or esters thereof.
11. The method according to claim 9, **characterized** in that the prostaglandin is 19R-hydroxy-PGE₂-methyl ester.
12. The method according to claim 9, **characterized** in that the prostaglandin is 19R-hydroxy-PGE₂- isopropyl ester.
13. The method according to any one of claims 9 – 12, **characterized** in that the prostaglandin is administered topically as a liniment, cream, ointment or gel or in solid form on the female genitals.

14. A pharmaceutical composition for the treatment of female sexual dysfunction, **characterized** in that said composition contains a prostaglandin being a selective EP₂ receptor agonist.

15. A pharmaceutical composition for the treatment of female sexual dysfunction, **characterized** in that said composition contains a prostaglandin chosen among 19-hydroxy-PGE₂, 19R-hydroxy-PGE₂, 18-hydroxy-PGE₂, 20-hydroxy-PGE₂, or salts or esters thereof.

16. A pharmaceutical composition for the treatment of female sexual dysfunction, **characterized** in that said composition contains 19R-hydroxy-PGE₂-methyl ester.

17. A pharmaceutical composition for the treatment of female sexual dysfunction, **characterized** in that said composition contains 19R-hydroxy-PGE₂- isopropyl ester.

HUMAN CORPUS C.

$n=8$




FIG. 1

% RELAXATION

RABBIT PENILE BLOOD VESSELS




FIG. 2

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 00/00006

A. CLASSIFICATION OF SUBJECT MATTER

IPC7: A61K 31/557

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Chemical Abstracts, Volume 87, No 7, 15 August 1977 (15.08.77), (Columbus, Ohio, USA), Spilman, C. H. et al, "Effects of 19-hydroxyprostaglandins on oviductal and uterine motility", page 58, THE ABSTRACT No 48352q, Prostaglandins 1977, 13 (4), 795-805	3-8
X	--	15-16
A	Chemical Abstracts, Volume 120, No 1, 3 January 1994 (03.01.94), (Columbus, Ohio, USA), Woodward, D.F. et al, "Identification of 19(R)-OH prostaglandin E2 as a selective prostanoid EP2-receptor agonist", page 161, THE ABSTRACT No 1279f, Prostaglandins 1993, 46 (4), 371-383	3-8
X	--	15-16

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

3 May 2000

Date of mailing of the international search report

09-05-1999

Name and mailing address of the ISA/

Swedish Patent Office

Box 5055, S-102 42 STOCKHOLM

Facsimile No. +46 8 666 02 86

Authorized officer

Göran Karlsson/EÖ

Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 00/00006

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 4164446 A (ARTHUR F. MARX ET AL), 14 August 1979 (14.08.79)	3-8
X	--	15
A	GB 1488006 A (ONO PHARMACEUTICAL CO. LTD.), 5 October 1977 (05.10.77)	3-8
X	--	15
P,X	WO 9902164 A1 (PHARMACIA & UPJOHN AB), 21 January 1999 (21.01.99)	15-17
P,A	WO 9921562 A1 (VIVUS, INC.), 6 May 1999 (06.05.99)	3-8
	-- -----	

INTERNATIONAL SEARCH REPORT

International application No.
PCT/SE00/00006**Box I** Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: **9-13**
because they relate to subject matter not required to be searched by this Authority, namely:
**A method for treatment of the human or animal body by therapy,
see rule 39.1.**
2. ☒ Claims Nos.: **1-2, 14**
because they relate to parts of the international application that do not comply with the prescribed requirements to such
an extent that no meaningful international search can be carried out, specifically:
see next sheet
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).:

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/SE00/00006

The expressions "selective EP2 receptor agonist or an active prostaglandin derivative thereof" in claim 1, "selective EP4 receptor agonist or an active prostaglandin derivative thereof" in claim 2 and "prostaglandin being a selective EP2 receptor agonist" in claim 14 are indefinite. According to PCT Article 6, the claims shall be clear and concise. The search has been aimed at the compounds given in claims 3-8.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/SE 00/00006

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4164446 A	14/08/79	AU 497219 B	07/12/78
		AU 7949875 A	30/09/76
		BE 827156 A	25/09/75
		CH 615159 A	15/01/80
		DE 2513222 A	02/10/75
		DK 125975 A	27/09/75
		FR 2265361 A,B	24/10/75
		GB 1501864 A	22/02/78
		IE 42704 B	08/10/80
		IT 1046092 B	30/06/80
		JP 50135283 A	27/10/75
		LU 72128 A	20/08/75
		NL 7503654 A	30/09/75
		SE 7503438 A	29/09/75
		US RE30287 E	27/05/80
		US 4054595 A	18/10/77
		US 4190670 A	26/02/80
		ZA 7501882 A	31/03/76
GB 1488006 A	05/10/77	AU 7810075 A	12/08/76
		BE 825459 A	12/08/75
		DE 2505519 A	14/08/75
		DK 50775 A	06/10/75
		FI 750371 A	14/08/75
		FR 2260335 A,B	05/09/75
		JP 50111050 A	01/09/75
		LU 71831 A	09/12/75
		NL 7501613 A	15/08/75
		SE 7501512 A	14/08/75
		US 4140712 A	20/02/79
		ZA 7500856 A	28/01/76
WO 9902164 A1	21/01/99	AU 8368298 A	08/02/99
		SE 9702681 D	00/00/00
WO 9921562 A1	06/05/99	AU 1125399 A	17/05/99
		US 5877216 A	02/03/99